



Baroreflex Sensitivity and Hemodynamic Recovery

Report prepared by

Eric A. Whitsel

Robert Angel

Rick O'Hara

Lixin Qu

Kathryn Carrier

Kathleen Mullan Harris



CAROLINA POPULATION CENTER | CAROLINA SQUARE - SUITE 210 | 123 WEST FRANKLIN STREET | CHAPEL HILL, NC 27516

Add Health is supported by grant P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 23 other federal agencies and foundations.

This document summarizes the rationale for and estimation of baroreflex sensitivity and hemodynamic recovery from the cardiovascular measures collected at the Wave V home exam. Whenever possible, data collection and methods in Wave V mirrored those of Wave IV to ensure comparability of data between waves. This document is one in a set of Wave V user guides. User guides are also available to describe protocols for the following biological measures at Wave V:

- Anthropometrics
- Cardiovascular Measures
- Medication Use – Home Exam
- Glucose Homeostasis
- Lipids
- Renal Function
- Hepatic Injury
- Inflammation and Immune Function
- Neurodegeneration

This research uses data from Add Health, a program project directed by Kathleen Mullan Harris and designed by J. Richard Udry, Peter S. Bearman, and Kathleen Mullan Harris at the University of North Carolina at Chapel Hill and funded by grant P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 23 other federal agencies and foundations. We gratefully acknowledge Carolyn Halpern, Nancy Dole, Joyce Tabor, and Sarah Dean for their dedication to the quality of the Add Health data in this document. Information on how to obtain the Add Health data files is available on the Add Health website (<https://addhealth.cpc.unc.edu/>).

Citations of this Add Health User Guide should use the following format:

Whitsel, EA, Angel R, O'Hara R, Qu L, Carrier K, Harris K. Add Health Wave V Documentation: Cardiovascular Measures Appendix I: Baroreflex Sensitivity & Hemodynamic Recovery 2024; Available from: <https://doi.org/10.17615/d4qt-5922>

Table of Contents

1. Introduction	3
2. Rationale	3
3. Estimation	3
3.1 Baroreflex Sensitivity (BRS).....	4
3.2. Hemodynamic Recovery	4
3.2.1 Pulse Rate Recovery (PRR)	4
3.2.2 Systolic Blood Pressure Recovery (SBPR)	4
3.3. Example Computation.....	4
4. Quality Control	5
4.1. Reliability	5
4.2. Validity	5
5. The Baroreflex Sensitivity Data File (bbrs5.xpt)	5
5.1. Structure	5
5.2. Contents.....	5
6. References	5

1. Introduction

This is an appendix to *Add Health Wave V Documentation: Cardiovascular Measures*¹. Please refer to that user guide for complete descriptions of the cardiovascular data collection procedures and measures disseminated by the study at that time. In addition to the measures described there, this appendix introduces three more constructed measures that are included in the Add Health Wave V data:

- Baroreflex sensitivity
- Pulse rate recovery
- Systolic blood pressure recovery

The rationale for their estimation and description of their quality control are provided below.

2. Rationale

The carotid sinus, aortic arch, heart, and lungs contain stretch receptors that are sensitive to increases in arterial, cardiac filling, and central venous blood pressure. These sinoaortic and cardiopulmonary baroreceptors project information about decreases in blood pressure to the spinal cord and brain which in turn, increase sympathetic and decrease parasympathetic stimulation of the heart and peripheral blood vessels. This baroreceptor reflex arc mediates the increase in pulse rate that usually accompanies decreases in systolic blood pressure over serial recordings, an autonomic phenomenon attributed to recovery from stress associated with instrumentation for sphygmomanometry and initial cuff inflation. Because baroreceptor reflexes prevent short-term fluctuation of blood pressure and their impairment is associated with cardiovascular diseases such as hypertension, diabetes, coronary heart disease, and heart failure^{2,3}, we estimated baroreflex sensitivity (BRS), pulse rate recovery (PRR) and systolic blood pressure recovery (SBPR) at Add Health Wave V.

3. Estimation

Estimation was restricted to 5,141 participants with non-missing sampling weights and all three resting, seated measures of SBP (mmHg) and PR (beats/min). It involved converting PR to its unit-corrected reciprocal, RR interval duration ($RR, ms = 60,000 \div PR$), and then running random-effects models implemented in Stata® 17 using MIXED and the Wave V Biomarker Weights⁴ to estimate BRS as the slope of the RR-SBP association, PRR as the increase in PR, and SBPR as the decrease in SBP across recordings.

3.1 Baroreflex Sensitivity (BRS)

The model used to estimate BRS (ms/mmHg) was given by: $RR_{ij} = \beta_0 + \beta_1 SBP_{ij} + \mu_{0j} + \mu_{1j} SBP_{ij} + e_{ij}$, where is the RR at the i^{th} measurement ($i = 1, 2, 3$) on the j^{th} participant ($j = 1, 2, 3 \dots, 5,141$), β_0 and μ_{0j} are the fixed and random intercepts, β_1 and μ_{1j} are the fixed and random slopes, and $BRS_j = \beta_1 + \mu_{1j}$.

3.2. Hemodynamic Recovery

3.2.1 Pulse Rate Recovery (PRR)

The model used to estimate PRR (beats/min) was given by: $PR_{ij} = \beta_0 + \beta_1 time_{ij} + \mu_{0j} + \mu_{1j} time_{ij} + e_{ij}$, where is the PR at the i^{th} measurement ($i = 1, 2, 3$) on the j^{th} participant ($j = 1, 2, 3 \dots, 5,141$), β_0 and μ_{0j} are the fixed and random intercepts, β_1 and μ_{1j} are the fixed and random slopes at times (0,1,2), and $PRR_j = \beta_1 + \mu_{1j}$.

3.2.2 Systolic Blood Pressure Recovery (SBPR)

The model used to estimate SBPR (mmHg) was given by: $SBP_{ij} = \beta_0 + \beta_1 time_{ij} + \mu_{0j} + \mu_{1j} time_{ij} + e_{ij}$, where is the SBP at the i^{th} measurement ($i = 1, 2, 3$) on the j^{th} participant ($j = 1, 2, 3 \dots, 5,141$), β_0 and μ_{0j} are the fixed and random intercepts, β_1 and μ_{1j} are the fixed and random slopes at times (0,1,2), and $SBPR_j = \beta_1 + \mu_{1j}$.

3.3. Example Computation

Exhibit 1 provides an example computation of $SBPR_j$ as the sum of the overall fixed slope and participant-specific, random slope (extracted using MIXED post-estimation commands). PRR_j and BRS_j were computed in an analogous way.

AID	Fixed Slope	Random Slope	SBPR
1	-1.114578	-0.267783	-1.382361
2	-1.114578	1.515851	0.401273
3	-1.114578	-0.340177	-1.454755
4	-1.114578	1.415925	0.301347
5	-1.114578	-1.246783	-2.361361

Exhibit 1. Example of estimating SBPR (mmHg)

4. Quality Control

4.1. Reliability

We assessed the short-term reliability of BRS, PRR and SBPR among a race/ethnicity- and sex-stratified random sample of 112 participants examined twice, 1-2 weeks apart among whom SBP and PR were measured following study protocol, typically by the same field interviewer and at approximately the same time of day. We used a nested, random-effects model to partition the variance of BRS, PRR and SBPR and estimated their reliability as the ratio of between-participant to total variance, i.e. a variance partitioning coefficient and 95% confidence interval.⁵ Greater than 80% of the variance in each of the three measures was attributable to that between participants. For BRS, the variance partitioning coefficient ranged from 0.91-0.96 over SBPs of 80-180 mmHg, and for PRR and SBPR, ranges were 0.91-0.91 and 0.83-0.86 over time.

4.2. Validity

See Cuthbertson, et al., 2014a and 2014b.^{6,7}

5. The Baroreflex Sensitivity Data File (bbrs5.xpt)

5.1. Structure

The structure of the disseminated baroreflex sensitivity data file is flat. This means that it is a respondent-level data file, where each respondent has one and only one record. The respondent's identifying number (the AID variable) will appear in the data file only once.

5.2. Contents

The baroreflex data file includes the variables below, which are described in the corresponding codebook documentation that also contains descriptive statistics.

<u>Variable Name</u>	<u>Variable Description</u>
AID	Respondent Identifier Number
H5BRS	Baroreflex sensitivity (ms/mmHg)
H5PRR	Pulse rate recovery (beats/min)
H5SBPR	Systolic blood pressure recovery (mmHg)

6. References

1. Whitsel, EA, Angel R, O'Hara R, Qu L, Carrier K, Harris K. (2020). *Add Health Wave V Documentation: Cardiovascular Measures*. UNC Chapel Hill: Carolina Population Center. Available from https://addhealth.cpc.unc.edu/wp-content/uploads/docs/user_guides/WaveVCardiovascularMeasuresUserGuide.pdf.

2. La Rovere MT, Pinna GD, Raczak G. (2008). Baroreflex Sensitivity: Measurement and Clinical Implications. *Ann Noninvasive Electrocardiol*;13(2):191-207.
3. Eckberg DL, Sleight P. (1992). *Human baroreflexes in health and disease*. New York: Oxford University Press.
4. Chen P, Harris KM. (2020). Construction of Wave V Biomarker Sample Weight. 2020. Carolina Population Center at the University of North Carolina at Chapel Hill. Available from <https://doi.org/10.17615/a5j6-3g23>.
5. Goldstein H. (1995). *Multilevel Statistical Models*. New York: Halsted Press.
6. Cuthbertson CC, Suchindran CM, Harris KM, Hussey JM, Halpern CT, Tabor JW, Williams RB, Killeya-Jones L, Whitsel EA. (2014a). Hemodynamic recovery and baroreflex sensitivity are reliable and associated with cardiovascular disease risk factors in a nationally representative sample of young US adults. *Circulation*;129(S1):AP3244.
7. Cuthbertson CC, Whitsel EA, Suchindran CM, Chen P, Cheng MM, Williams RB, Tabor JW, Killeya-Jones LA, Hussey JM, Halpern CT, Harris KM. (2014b). Add Health Wave IV Documentation. Cardiovascular Measures Appendix I: Baroreflex Sensitivity & Hemodynamic Recovery. Available from https://addhealth.cpc.unc.edu/wp-content/uploads/docs/user_guides/Baroreflex_Sensitivity.pdf.